

R_2 is C_{1-6} alkyl;

R_3 is selected from hydrogen, C_{1-6} alkyl, mono- or di(C_{3-6} cycloalkyl)methyl, C_{3-6} cycloalkyl; C_{3-6} alkenyl; hydroxy C_{1-6} alkyl, C_{1-6} alkylcarbonyloxy C_{1-6} alkyl and C_{1-6} alkyloxy C_{1-6} alkyl;

R_4 and R_5 are independently selected from C_{1-8} alkyl, mono- or di(C_{3-6} cycloalkyl)methyl, Ar^1CH_2 , C_{3-6} alkenyl, C_{1-6} alkyloxy C_{1-6} alkyl, hydroxy C_{1-6} alkyl, thienylmethyl, furanymethyl, C_{1-6} alkylthio C_{1-6} alkyl, morpholinyl, mono- or di(C_{1-6} alkyl)amino C_{1-6} alkyl, di(C_{1-6} alkyl)amino, C_{1-6} alkylcarbonyl C_{1-6} alkyl, C_{1-6} alkyl substituted with imidazolyl; or a radical of the formula $-(C_{1-6}alkanediy)-O-CO-Ar^1$;

or R_3 and R_4 taken together with the nitrogen atom to which they are attached form a pyrrolidinyl, piperidinyl, homopiperidinyl or morpholinyl group, optionally substituted with C_{1-6} alkyl or C_{1-6} alkyloxy;

Ar is selected from phenyl substituted with 1, 2 or 3 substituents independently selected from halo, C_{1-6} alkyl, trifluoromethyl, cyano, C_{1-6} alkyloxy, benzyloxy, C_{1-6} alkylthio, nitro, amino and mono- and di(C_{1-6} alkyl)amino; and pyridinyl substituted with 1, 2 or 3 substituents independently selected from halo, C_{1-6} alkyl, trifluoromethyl, hydroxy, cyano, C_{1-6} alkyloxy, benzyloxy, C_{1-6} alkylthio, nitro, amino, mono- and di(C_{1-6} alkyl)amino and piperidinyl; and

Ar^1 is selected from phenyl, pyridinyl, and phenyl substituted with 1, 2 or 3 substituents independently selected from halo, C_{1-6} alkyl, C_{1-6} alkyloxy, di(C_{1-6} alkyl)amino C_{1-6} alkyl, trifluoromethyl and C_{1-6} alkyl substituted with morpholinyl.

Please add the following claims:

19. (new) The method of claim 5 wherein said disorder is anxiety or depression.
20. (new) The method of claim 5 wherein said disorder is irritable bowel syndrome, inflammation, anorexia nervosa, seizures, or substance abuse.